# EFFECTS OF DISTURBANCE ON KILLER WHALES

David Bain,
University of Washington

#### Introduction

**Sources of Disturbance Mechanisms of Impact** 

#### Methods

Observation Experimentation

#### Results

Behavior States
Surface Active Behavior
Movement Patterns
Effects of mid-frequency sonar

#### **Discussion**

Zones of Influence
Effects on foraging efficiency
Population Scale Effects
Management Options

### Sources of Disturbance

- Explosives
- Airguns
- Mid-Frequency Sonar
- LFA Sonar
- Acoustic Tomography
- Industrial
- Acoustic Harassment/Deterrent Devices

- Tourist Vessels
- Commercial Fishing
- Freight Traffic
- Aircraft
- Biological
- Physical

### Mechanisms

- Collisions with vessels
- Pollution
- Noise
- Physical presence
- Stress

### Noise Impact Mechanisms

- Masking
- Threshold Shifts
- Displacement
  - Excess Energy Expenditure
  - Impaired Foraging Efficiency
- Behavioral Changes
- Effects on Prey

### Observations of focal whales

- land-based
- theodolite (objective and repeatable)
- measure behavior with no boats around
- compare to behavior with different levels of boat activity (number, proximity, operating practices)



### Experiments on focal whales



- observe a whale for 20min without boats
- approach focal with experimental boat
- parallel the whale at 100m for 20min
- compare pre-exposure, exposure, and postexposure data

### Sampling all whales, all activities


15min scans – 8yr

no focal bias


 Markov-chain modeling

 whales likelier to stop feeding after
 15min with boats present than absent


 not all activities equally vulnerable to disturbance




### Effect of boats on activity budgets



## Surfacing Patterns






·· whale

boat


### So what?



- not all behaviors changed
- results consistent with horizontal avoidance
- swim 13% farther to get where they need to go
- that may cost energy

Williams et al. 2002a

### **Speed Matters**




- whales used the same, but more obvious, response as to a paralleling zodiac
- response to leapfrogging at ~150m was detectable with observations on only 10 animals

(Williams et al. 2002b)

#### **Distance Matters**

Change in Probability of Feeding as a Function of Proximity to the Closest Boat



## **Distance Matters** (Southern Residents 2003)


DISTANCE	<400	>400
Dive Time	<b>56.2</b>	44.2
Speed	6174	6462
Deviation	29.3	23.4
Directness	74.2	77.8
<b>Surface Active</b>	0.658	2.857

#### significantly different


non-significant but in same direction as Northern Residents

#### **Distance Matters**

Surface Active Behavior as a Function of Observing Distance (SR 2003)



## Why is leapfrogging so disturbing?




- when an outboard speeds up, noise gets louder, and higher in frequency
- noise placed directly in front of whale, which causes most masking

received noise of fast boat at 500m equaled slow boat at 100m


Deep Water Path Established
Tail Slap

Distance = 47 km, 26 nm Time: 1047




No Deep Water Path
J Pod behaving normally

Distance = 47-22 km, 26-12 nm Time: 1047-1134




## Deep water path established J Pod turns to move away

Distance = 22 km, 12 nm Time: 1134




## Deep Water Path J Pod behaving abnormally Distance = 18-22 km, 10-12 nm Time: 1134-1314




No Deep Water Path J Pod normal (slow, spread) Time: 1314-1355

**Distance** = 15-22 km, 8-12 nm




#### Deep Water Path J Pod Abnormal

Distance = 13-15 km, 7-8 nm Time: 1355-1407




No Deep Water Path J Pod staying in shadow zone Time: 1407-1432

**Distance = 4-13 km, 2-7 nm** 




## **Closest approach J Pod splitting**

**Distance = 3 km, 1.7 nm Time: 1432** 




## Shoup moving away J Pod splitting, part moving away Time: 1432-1438





Sonar off


Distance = 4+ km, 2+ nm


J Pod split, moving offshore, spreading out Time: 1438-1452






















## Foraging tactics




#### Relative Prey Availability Due to Noise-Induced Threshold Changes




## **Effects of Population Size and Cumulative Effect Size on Population Growth**

Effect of Whale Watching on Population Growth Rate (Shape Parameter = 11.3)



- negligible effects expected well below carrying capacity
- disturbance costly near carrying capacity

#### Correlation ( $r^2=0.42$ , p < .01) of Fleet Size with Population Dynamics



- Role in Decline
  - Could account for much of recent decline in Southern Residents
- Role in Recovery
  - Toxins would slow recovery in Southern Residents
  - Recovery of fish stocks could offset effects of whale watching

(Bain et al.)

# Reducing the Cumulative Effect of Whale Watching

Total Quota Based on Potential Biological Removal

Limited Entry/Individual Transferable Quotas

License Fee to Cover Management Costs

Time and area closures

Changing operating practices to reduce impact

Quieter Vessels

Increasing Viewing Distance

Slowing Down Near Whales

#### Thanks to colleagues

Sue Kruse, Dave Briggs Janice Waite Andrew Tr Marilyn Dahlheim, Bob Otis, Rob Williams, Rich Osborne, Jodi Smith, Patrick Miller, John Ford Carlos Alvarez-Flores, Glenn VanBlaricom, and many others

#### Thanks for financial and logistical support

**National Science Foundation University of California, Santa Cruz Stubbs Island Charters** Joseph M. Long **Data General Corporation** National Marine Fisheries Service United States Geological Sur Minerals Management Service Orca Free **Orca Conservancy Orca Relief Citizens Alliance Friday Harbor Laboratories** Whale Watch Operators Association NW **NSERC** Weber Fund **WDCS** TRFF **IFAW** Bion **BC Parks** 

**DFO**